

Species

Partial replacement of fish meal with mangrove based plant ingredients and its effect on water quality, growth performance and length-weight relationship of freshwater prawn *Macrobrachium rosenbergii*

Mondal K^{1,*}, Ghosh R², Bhattacharyya SB³, Zaman S⁴, Mallik A⁵, Das M⁶, Mitra A⁷

1. Research Scholar, Department of Marine Science, University of Calcutta, 35 B.C. Road, Kolkata – 700019, India
2. Senior Research Scholar, Department of Marine Science, University of Calcutta, 35 B.C. Road, Kolkata – 700019, India
3. Research Associate (III), Department of Marine Science, University of Calcutta, 35 B.C. Road, Kolkata – 700019, India
4. DST Women Scientist, Department of Marine Science, University of Calcutta, 35 B.C. Road, Kolkata – 700019, India
5. Research Scholar, Division of Molecular Medicine, Bose Institute, P 1/12, C. I. T. Road, Scheme – VIIM, Kolkata – 700054, India
6. Professor, Department of Zoology, University of Calcutta, 35 B.C. Road, Kolkata – 700019, India
7. Associate Professor, Department of Marine Science, University of Calcutta, 35 B.C. Road, Kolkata – 700019, India

*Corresponding author: Research Scholar, Department of Marine Science, University of Calcutta, 35 B.C. Road, Kolkata – 700019, India, Mail: bottu.marine@gmail.com, Mobile No. 09007794059.

Received 13 March; accepted 22 April; published online 01 May; printed 16 May 2013

ABSTRACT

Partial replacement of 30% fish meal was done with 10% plant ingredients comprised of macroalgae (*Enteromorpha intestinalis*) and saltmarsh grass (*Porteresia coarctata*) to prepare freshwater prawn (*Macrobrachium rosenbergii*) feed. The floral based feed was applied to culture *M. rosenbergii* in Indian Sundarbans region. The water quality parameters (temperature, salinity, pH, D.O, nutrient concentrations, chlorophyll a, transparency, soil organic carbon) and growth performance (average initial weight, average final weight, DGR, SGR, survival rate, FCR, total yield) were monitored throughout the culture period. The pond health was best where diet-3 was applied compared to the other ponds. The condition state was also monitored based on length-weight relationship (LWR). The growth was uniform and maximum in cultured species fed with diet-3 compared to that of other formulated diets (diet-C, diet-1 and diet-2). $LWR (W = 0.008 \times TL^{2.9486})$ indicated isometric growth of *M. rosenbergii* fed with diet-3. The trial experiment thus demonstrates the efficacy of diet-3 (*P. coarctata* based feed) in maintaining the overall growth performance and condition state of the cultured species as well as the environmental health of the culture pond.

Keywords: Fish meal, *E. intestinalis*, *P. coarctata*, water quality, growth performance, length-weight relationship, *M. rosenbergii*.

Abbreviation: DGR-Daily Growth Rate, SGR-Specific growth rate, FCR-Feed conversion ratio, LWR-Length weight relationship, DO-Dissolved oxygen.

1. INTRODUCTION

Freshwater prawn (*M. rosenbergii*) is one of the most commercially important species in the fisheries sector of India as a source of animal protein and foreign exchange earner. The species is widely distributed in freshwater as well as brackishwater bodies like ponds, rivers, canals and estuaries (Ahmed, 1957). The species is the largest of the freshwater prawns and has been classified as a benthophagic omnivore based on analysis of foregut contents (Weindenbach, 1982; Cohen and Ra'anani, 1983). It has recently been identified as an important shellfish species for culture in South Asian countries after significant losses of penaeid shrimp culture in mid-1990s due to viral diseases (Paez-Osuna, 2001; Moss et al., 2001). In India, 36,640 hectares are under scampi culture with an annual production of about 39,000 tons from monoculture and as one of the components in polyculture (Lokare and Anis, 2000). The scampi are fed with farm-made or commercial feed in order to boost up the production volume. While

prawns can receive substantial nutritional benefit from natural food at relatively low biomass densities of less than 1000 kg ha⁻¹ (Tidwell et al., 1997), at higher stocking densities individuals may be more dependent on prepared diets (Tidwell et al., 1999). For achieving successful production in prawn culture, a great deal of consideration is generally given to feeding and management. Feed costs constitute 40-60% of operational costs in production of the freshwater prawn (D'Abramo and Sheen, 1991). Fish meal makes up a major part of the formulated feed for many carnivorous fish species as a protein source. However, the recent scarcity, rising cost and uncertain consistency in supply of fish meal have increased the efforts to search for the alternative protein sources to fish meal in aquaculture by the fish nutritionists and feed manufacturers. For the purpose of producing low cost effective diets, high protein feedstuffs for fish meal substitution have been investigated throughout the world (Hasanuzzaman et al., 2009).

Indian Sundarbans (world's largest mangrove delta, a World Heritage Site and a Biosphere Reserve) located in the lower Gangetic region of the Indian sub-continent at the apex of Bay of Bengal (within latitude 21°13' N to 22°40' N and longitude 88°03' E to 89°07' E) offers a congenial environment in terms of salinity and other hydrological parameters for the growth and culture of scampi (Mitra et al., 2006). The deltaic complex sustains some 34 true mangrove species and 65 mangrove associate floral species (Mitra,

Fish meal:

It is a commercial product made from fish and the bones and offal from processed fish. It is a brown powder or cake obtained by drying the fish or fish trimmings, often after cooking, and then grinding it. Fishmeal is a nutrient-rich and high protein supplement feed ingredient that stores well, and is used primarily in diets for domestic animals and sometimes as a high-quality organic fertilizer.

Performance of *M. rosenbergii*

The present programme is a comparative account of the growth performance of freshwater prawn (*M. rosenbergii*) due to use of different categories of formulated fish feed. The comparison reflects the efficacy of the feed prepared from *Porteresia coarctata* in boosting the prawn growth and maintaining the aquatic health.

Mondal et al.

Partial replacement of fish meal with mangrove based plant ingredients and its effect on water quality, growth performance and length-weight relationship of freshwater prawn *Macrobrachium rosenbergii*, Species, 2013, 3(8), 15-21,

© The Author(s) 2013. Open Access. This article is licensed under a [Creative Commons Attribution License 4.0 \(CC BY 4.0\)](http://creativecommons.org/licenses/by/4.0/).

Table 1 Ingredients (%) of control diet (diet-C) and experimental diets (diet-1, diet-2 and diet-3)

Ingredients	Diet- C	Diet-1	Diet-2	Diet-3
Soybean meal	10	10	10	10
Mustard oil cake	34	34	34	34
Rice bran	17	17	17	17
Wheat bran	7	7	7	7
Vitamin + mineral mixture	2	2	2	2
Fish meal	30	20	20	20
Powder of <i>Enteromorpha intestinalis</i>	-	10	5	-
Powder of <i>Porteresia coarctata</i>	-	-	5	10

Table 2 Proximate composition of control and experimental diets

Proximate composition (%)	Diet-C	Diet-1	Diet-2	Diet-3
Crude protein	33.47 ± 1.2	34.94 ± 1.2	34.93 ± 1.8	36.49 ± 1.3
Soluble carbohydrate	34.12 ± 1.6	33.89 ± 1.9	35.01 ± 2.5	36.18 ± 0.9
Crude fat	6.1 ± 0.52	6.1 ± 0.99	5.8 ± 0.46	6.2 ± 0.23
Ash	11.4 ± 0.21	10.2 ± 0.12	11.9 ± 0.31	12.1 ± 0.23
Moisture	10.4 ± 0.22	9.5 ± 0.11	10.1 ± 0.34	10.2 ± 0.99

Macroalgae:
These are large aquatic photosynthetic plants that can be seen without the aid of a microscope. On the basis of the pigment present, macroalgae can be of different colors like green, red, brown and blue. The most familiar types can be generally divided into three groups: Chlorophyta, Rhodophyta and Phaeophyta.



Figure 1
Live and healthy macroalgae (*Enteromorpha intestinalis*)

Saltmarsh grass:
A saltmarsh grass is a vital biotic component of the marshy ecosystem found along the coastal and deltaic zones. They are highly productive floral species and have wide range of salinity tolerance. In the present study, the saltmarsh grass *Porteresia coarctata* is the marsh grass species that has been used as one of the ingredient for prawn feed preparation. Apart from its nutritional value, the species has ecological value as it brings stability to the newly formed island. It is also referred to as pioneer species in the process of ecological succession.



Figure 2
Saltmarsh grass (*Porteresia coarctata*) vegetation on intertidal mudflat

2000). A considerable percentage (73%) of people living in the deltaic region have opted pisciculture as one of the important livelihoods. However, there is scarce concept of scientific feed preparation and water management in the region due to which the aquaculture sector often gets devastated with disease problems. Deterioration of pond water quality due to inclusion of trash fish in the diet of cultured species is also an associated problem with the

prawn culture system in this region (Mitra, 1994). Finding alternative, traceable, high quality protein to fishmeal is a key area of organic aqua feed research. Research into organic formulations has been steadily increasing over the last several years.

On this background, the present study aims to evaluate the effects of partially replacing the ingredients of animal origin of conventional prawn feed with macroalgae (*E. intestinalis*) and saltmarsh grass (*P. coarctata*) powder. The protein levels of *Enteromorpha* spp. generally range between 5-20% of dry weight and is abundantly available in saline water bodies of Sundarbans (Mitra and Banerjee, 2006). Because of high protein content, protein concentrates (PC's) of seaweeds have become more important for the food industry, especially in developed countries (Wong and Cheung, 2001). The recent utilization of macroalgae as shrimp feed is also gaining momentum in deltaic ecosystem of Indian Sundarbans (Banerjee et al., 2010). The salt marsh grass, *P. coarctata* is another cheapest source of protein (18%), widely available in the intertidal mudflats of Sundarbans (Mitra and Banerjee, 2006).

2. MATERIALS AND METHODOLOGY

2.1. Collection and processing of macroalgae and saltmarsh grass

Live and healthy *E. intestinalis* (Figure 1) and *P. coarctata* (Figure 2) were collected from the intertidal mudflat in the mangrove belt of Kakdwip region (21°52'35.7" N latitude and 88°11' 55.0" E longitude) during low tide condition and washed thoroughly with double distilled water to remove epiphytes and other extraneous matter. Collected materials were sun dried and ground to make a fine powder. Three categories of powder were prepared:
Category 1- *E. intestinalis* powder
Category 2 - *P. coarctata* powder
Category 3 - Equal mixture (1:1) of *E. intestinalis* and *P. coarctata* powder

2.2. Diet preparation

Four different types of diets were formulated viz. control diet (diet-C) and three experimental diets (diet-1, diet-2, and diet-3). The major source of protein in diet-C was from the fish meal. The fish meal protein was partially replaced in the experimental diets at a rate of 10% of feed with the powder of *E.intestinalis*, *P.coarctata* and mixture (1:1) of both powders. The diet-C was provided to pond-1 (treated as control) and subsequently diets-1, 2, 3 were provided to ponds-2, 3 and 4 respectively (treated as experimental). All the ponds had three replicates each. Formulation of control and experimental diets are presented in Table 1. The proximate chemical composition of the diets after preparation was done according to AOAC (1980) as provided in Table 2.

2.3. Experimental prawn farming

The eight month culture experiment of *M. rosenbergii* was conducted during February to September' 2011 in farm condition at Kakdwip region of Indian Sundarbans (Figure 3). Twelve ponds of different size were selected for trial of

Table 3 Effect of different diets on hydrological characteristics and soil organic carbon of culture ponds

Parameters	Pond-1 (provided with diet-C)	Pond-2 (provided with diet-1)	Pond-3 (provided with diet-2)	Pond-4 (provided with diet-3)
Temperature (°C)	32.7±0.29 ^a	32.76±0.15 ^a	32.7±0.13 ^a	32.7±0.12 ^a
Salinity (psu)	0.56±0.38 ^a	0.59±0.42 ^a	0.41±0.44 ^a	0.26±0.43 ^a
pH	7.19±0.31 ^b	7.62±0.04 ^a	7.84±0.03 ^a	7.81±0.01 ^a
DO (mg l ⁻¹)	4.32±0.48 ^c	5.06±0.11 ^b	5.25±0.15 ^{ab}	5.35±0.18 ^a
Nitrate (µgat l ⁻¹)	33.5±3.45 ^a	27.0±3.07 ^b	25.2±3.54 ^{bc}	23.3±4.34 ^c
Phosphate (µgat l ⁻¹)	4.06±1.22 ^a	3.43±0.49 ^b	3.35±0.52 ^c	2.76±0.79 ^c
Silicate (µgat l ⁻¹)	59.2±6.20 ^a	48.1±3.40 ^b	43.1±4.16 ^c	45.2±5.47 ^{bc}
Chlorophyll a (mg m ⁻³)	3.10±0.43 ^b	2.89±0.31 ^a	2.92±0.41 ^a	3.09±0.43 ^b
Transparency (cm)	18.3±2.52 ^b	19.8±2.83 ^b	25.7±1.37 ^a	23.3±2.45 ^a
Soil organic carbon (%)	4.73±1.15 ^a	3.50±1.36 ^b	3.19±0.47 ^b	3.06±0.78 ^b

The results are means ± S.D. (standard deviation) and the figures with same superscripts in the same row are not significantly different (p < 0.05).

Table 4 Growth performance of *M. rosenbergii* fed with mangrove based diets

Parameters	Diet-C	Diet-1	Diet-2	Diet-3
Initial average weight (gm)	0.02±0.01 ^a	0.02±0.01 ^a	0.02±0.01 ^a	0.02±0.01 ^a
Final average weight (gm)	86.26±2.68 ^c	80.83±1.31 ^d	90.53±3.25 ^b	92.87±2.1 ^a
Daily growth rate (gm day ⁻¹)	0.37±0.01 ^b	0.33±0.01 ^d	0.37±0.01 ^c	0.38±0.20 ^a
Specific growth rate (% day ⁻¹)	3.57±0.01 ^c	3.45±0.05 ^d	3.66±0.02 ^b	3.69±0.01 ^a
Survival rate (%)	63.9±2.36 ^d	66.3±2.53 ^c	73.7±3.53 ^b	79.0±3.00 ^a
Feed Conversion Ratio	2.02±0.02 ^a	1.89±0.01 ^b	1.76±0.01 ^c	1.68±0.02 ^d
Total yield (kg ha ⁻¹)	1102 ^c	1071 ^d	1334 ^b	1467 ^a

The results are means ± S.D (standard deviation) and the figures with same superscripts in the same row are not significantly different (p < 0.05).




Figure 3 Map showing the location of culture site at Kankdip region of Indian Sundarbans

experimental and control diets in triplicate. Ponds were prepared following standard procedure. Seeds were procured from local hatchery and stocked at a rate of 2 individual/m² with average initial length 1.0 cm and 0.02 gm initial average body weight in each pond. The mean stocking weight was determined from a sample of 100 prawn seeds that were blotted to free from water. Before stocking all the prawn seeds are well acclimatized to avoid temperature and pH shocks (Sarver et al., 1982). Prawns were fed to satiation twice a day. The feeding levels were adjusted by checking the feeding trays two hours after serving the diets. Prawns were harvested at the end of 240 days. Total body length, weight and number of prawns from each pond were recorded.

2.4. Growth performance

The following growth parameters were measured for the cultured species from the control and experimental ponds:

- Average initial weight and final weight were measured using pan balance.

- Daily growth rate (DGR) was calculated using the expression: DGR (gm day⁻¹)= W_f (final wet wt) - W_i (initial wet wt)/ t (in days)
- Specific growth rate (SGR) was calculated after the harvesting of prawns as per the expression: SGR (%) = ln (final weight) – ln (Initial weight) / days of experiment x 100 (De Silva and Anderson, 1995).
- Feed Conversion Ratio (FCR) was calculated after the harvesting of prawns as per the expression: FCR = feed consumed (dry weight) / live weight gain (wet weight), (De Silva and Anderson, 1995).
- Survival rate was calculated as per the formula: Survival rate = Final no. of prawns / initial no. of prawns x 100 (De Silva and Anderson, 1995).
- Length-weight relationship (LWR): The biometric measurements of the cultured species were made with graduated scale. The measurements taken were total length (TL) from the tip of the rostrum up to the tip of the telson. The measurements were made to centimeter (cm) as described by FAO species identification sheets for fishery purposes (Fischer et al., 1981). The specimens were also weighed using a pan balance for taking the total weight (W). The mathematical relationship between length and weight was calculated using the conventional formula (Ricker, 1973):

W= a. TL^b

Where, W= Total weight
TL= Total length
a = Proportionality constant
b = Isometric exponent

2.5. Water quality parameters

Water quality parameters such as temperature, pH, salinity, dissolved oxygen (DO), transparency and nutrient concentrations (nitrate, phosphate and silicate) for all the ponds were measured on monthly basis following standard procedures as stated in Strickland and Parsons (1972). Chlorophyll a concentration was measured spectrophotometrically after acetone extraction as per APHA (1995) and soil organic carbon was analyzed by Walkey and Black method (1934).

2.6. Statistical Analysis

The data generated for all biotic and abiotic parameters were expressed as mean ± S.D. Significance of variations in the water quality, growth performance and length-weight relationship of the cultured species was tested by using one way Analysis of Variance (ANOVA) which was followed by Duncan's Multiple Range Test for significant values using SPSS software. Values were considered significant at 5% level of significance (Gomez and Gomez, 1984).

3. RESULTS

3.1. Water Quality

The water quality of the culture ponds are given in Table 3. During the experimental period, temperature varied in the range 32.70 to 32.76°C, pH varied in the range of 7.19 to 7.84 and dissolved oxygen (DO) in the range 4.32 to 5.35 mg l⁻¹. Salinity varied in the range between 0.26 to 0.59 psu. All the water parameters were found to be in the range of tolerance limit of *M. rosenbergii* (New and Singholka, 1985). In general, nutrient concentrations (nitrate, phosphate and silicate) increased as culture proceeded. Nitrate concentration was found to be highest in pond-1 (33.5 µg-at l⁻¹) and lowest in pond-4 (23.3 µg-at l⁻¹). Similar trends were observed for phosphate and silicate concentrations. The indicator of primary productivity i.e. chlorophyll a concentration also increased with time in all the culture ponds. Highest chlorophyll a level was observed in pond-1 (3.1 mg m⁻³) and lowest in pond-2 (2.89 mg m⁻³). Organic carbon content was highest in pond-1 showing a value of 4.73%, where fish meal was used as feed ingredient. A low organic carbon value (3.06%), an indication of better pond environment is observed in pond-4 applied with diet-3. Higher water transparency was recorded from pond-3 (25.7 cm) and pond-4 (23.3 cm) respectively.

3.2. Growth performance of freshwater prawn

Growth data of the experiment are given in Table 4. The lowest survival rate of prawn was observed in pond-1 fed with diet- C (63.9%) and the highest was observed in pond-4 fed with diet-3 (79.0%). The average final weight of the prawn varied from 80.83 to 92.87 g. The prawns fed with

diet-3 showed the highest average weight throughout the 240 days of feeding trial (Figure 4). The prawns that fed on diet-3 and diet-2 gained significantly higher average weight compared to the prawns fed on diet- C and diet-1 ($p < 0.05$). SGR for the cultured species varied between 3.45 and 3.69 % day⁻¹. It was evident that SGR was lowest in case of diet-1 and the highest in prawns fed with diet-3. The lowest production of prawn (1071 kg ha⁻¹) in the pond condition was obtained where diet-1 was applied and the highest was obtained in case of prawns treated with diet-3 (1467 kg ha⁻¹). The FCR values of the different types of experimental diets varied between 1.68 to 2.02. The highest FCR was observed in pond-1 (provided with diet-C) and the lowest in pond-4 (provided with diet-3). DGR for the cultured species varied between 0.33 to 0.38 gm day⁻¹ with highest value in pond-4 (0.38 gm day⁻¹) and lowest in pond-2 (0.33 gm day⁻¹).

3.3. Length-weight relationship

The mathematical relationship between length and weight of *M. rosenbergii* fed with diet-C and diets- 1, 2 and 3 are shown in figures 5 to 8. Isometric exponent (b) values ranged between 2.71 to 2.95. Lowest and highest values were observed in diet-1 and diet-3 treated ponds respectively. In the present study, the isometric exponent value (b=2.95) observed in *M. rosenbergii* fed with diet-3 is an indication of isometric growth.

4. DISCUSSION

4.1. Water Quality

Prawn growth and health are influenced by the pond bottom soil (sediment) and water quality parameters. Therefore,




Figure 4

Growth pattern of *M. rosenbergii* fed on different mangrove based diets




Figure 5

Length-weight relationship of *M. rosenbergii* fed with control diet (diet-C)




Figure 6

Length-weight relationship of *M. rosenbergii* fed with diet-1 (prepared with *E. intestinalis*)




Figure 7

Length-weight relationship of *M. rosenbergii* fed with diet-2 (prepared with equal mixture of *E. intestinalis* and *P. coarctata*)




Figure 8

Length-weight relationship of *M. rosenbergii* fed with diet-3 (prepared with *P. coarctata*)

management of pond sediments and water quality is the key for successful aquaculture practice. Therefore, properties of the pond sediment and processes occurring at the bottom soil and in the soil-water interface determine the water quality. During the study period, all the hydrological parameters were found to be within the acceptable limits for culturing *M. rosenbergii* as reported in various literatures (Daniel, 1981; Sandifer and Smith, 1985; Boyd, 1990; New, 1995; D'Abramo and Brunson, 1996; Hall and Van Hamm, 1998). The experimental treatments showed better aquatic environment compared to the control treatment where conventional diet (included with fish meal) was provided. Surface water temperature in all the culture ponds had more or less parallel trend of variation throughout the study period which is reflected in the ANOVA results. This uniformity in temperature profile is due to the location of culture ponds in the same site that experience similar weather and climate. Similar observations were recorded for salinity profile which may be due to the same water source of Kalnagini river of Hugli-Matla estuarine complex. No significant variation was observed for pH due to the traditional practice of liming in lower gangetic region at regular interval of time. D.O is an important component regulating the aquatic life. Its level varied significantly between ponds ($p < 0.05$) as per the order pond 4>pond 3>pond 2>pond 1. This clearly shows the role of mangrove based feed in upgrading the water quality of the culture ponds which may be attributed to the minimum deposition of organic matter at the bottom of the pond. The better aquatic environment in the experimental treatments was reflected through lowering of nutrient levels and subsequent increment of phytoplankton (chlorophyll *a*) level. ANOVA results confirmed significant variation ($p < 0.05$) in nutrient load between the treatments which may be attributed to leaching of feed ingredients (particularly animal component of the feed) and also the faecal matter that generates ammonia (Mitra and Choudhury, 1995). The increased water transparency in case of ponds 2, 3 and 4 may be due to the unique binding properties of the selected mangrove based plant ingredients used to prepare the feed thereby reducing the suspended particulate matter. Similar observations were recorded during culture of *Penaeus monodon* in Sundarbans through application of formulated feed prepared from the red algae *Catenella repens* (Banerjee et al., 2010). The organic carbon content at the pond bottom soil exhibited a decreasing trend with time in case of experimental treatments compared to the control, thus reflecting the efficient utilization of mangrove based feed by the cultured species.

4.2. Growth performance of freshwater prawn *M. rosenbergii*

The present study confirms significant variation in survival rate, DGR and SGR of the cultured species between the control and experimental treatments ($p < 0.05$) which may be

attributed to differences in the component and proportions of feed ingredients. The growth pattern of *M. rosenbergii* shows uniformity up to 75 days in all the culture ponds. From 75th day onwards there is an exponential growth in the species fed with mangrove based diet. Prawns fed with diet-3 (where 30% fish meal is partially replaced by 10% *P. coarctata* powder) showed maximum average weight followed by diet-2 (where 30% fish meal is partially replaced by equal proportion of *P. coarctata* & *E. intestinalis* powder), diet-C (comprised of 30% fish meal) and diet-1 (where 30% fish meal is partially replaced by 10% *E. intestinalis* powder). The FCR and SGR followed the order diet-C>diet-1>diet-2>diet-3 and diet-3>diet-2>diet-C>diet-1 respectively indicating the efficacy of diet-3 in maintaining the pond environment and its utilization for growth of the cultured species. Significant differences in FCR were observed among the experimental treatments ($p < 0.05$). The FCR values observed for diet-1, diet-2 and diet-3 during the study period are lower than the values (2.18-2.43) reported by William, D'Abramo, Fondren and Durant (1995) and Hossain et al., (2000) for *M. rosenbergii*. Even higher FCR values were observed (3.7-5.6) for *M. rosenbergii* cultured in concrete tanks fed with formulated feed containing 34% protein (Siddiqui et al., 1997). The highest survival rate was recorded from the prawns fed with diet-3 due to the suitable environment of the particular culture system. The unique growth performance in the present study with mangrove based feed is in accordance with the findings of Weidenbach (1980) who reported that prawns are able to adjust with the absence of feed pellets by increasing consumption of available vegetation. According to Tidwell et al., (1995), prawns are able to adjust to reductions in the nutritional value of prepared diets (i.e. protein source and vitamin and mineral content) by increasing predation on natural fauna (i.e. macro invertebrates) in the culture system. The highest production (1467 kg ha⁻¹) of prawns in 8 month culture period was obtained in diet-3. This is because of higher survival and weight gain observed in this treatment (Hossain and Paul, 2007). Digestibility studies in *M. rosenbergii* have indicated that the species can efficiently digest both plant and animal protein sources (Ashmore et al., 1985). The overall growth performance of *M. rosenbergii* fed with diet-3 might be attributed to better nutritive balance of the ingredients especially essential amino acids (Hossain and Paul, 2007). May be the utilization of mangrove based plant ingredients in different percentage in feed have been found to be more effective than a single floral source as ingredients of fish feed (Hossain and Jauncey, 1990).

4.3. Length-weight relationship

Length-weight relationship study is an important index to measure the variation in the growth of individual prawn or group of prawns (Jayachandran and Joseph, 1988) as well as for stock assessment (Abowweyere and Williams, 2008). It also plays a significant role in studying the growth, rate of feeding, metamorphosis, fatness, onset of maturity, gonadal development and general well-being of the fish population (Le Cren, 1951; Pauly, 1993). Variation in the length-weight relationship obtained between the experimental and control treatments for *M. rosenbergii* often arises due to the variation in food composition and environmental conditions as revealed in earlier studies (Abrahamson, 1966; Kuris et al., 1987). Earlier fish biologists critically analyzed the significance of LWR and stated that when 'b' is equal to 3 then growth is isometric Enin (1994) and when it is less than or greater than 3 it is allometric (Wootton, 1992). In the present study, cultured species fed with diet-3 exhibited highest b-value of 2.95 which is proximal to the standard isometric value $b = 3$. Based on *t*-test ($t = 0.8097$, $p < 0.05$), no significant variation is observed between the standard isometric value and the *b*-value obtained in the present study (Biswas et al., 2011) thus indicating a proportionate increase of length and weight. The isometric exponent (*b*) values for diet-C, diet-1 and diet-2 differed significantly from diet-3 ($p < 0.05$) which indicates a disproportionate length-weight relationship in cultured species (allometric growth) fed with diet-C (where $b = 2.71$), diet-1 (where $b = 2.73$) and diet-2 (where $b = 2.84$) respectively.

5. CONCLUSION

The deltaic complex of Sundarbans in the lower Gangetic region offers a unique opportunity for the culture of *M. rosenbergii*. However the endeavor mostly becomes unsuccessful due to the use of traditional fish feed in the region. The traditional feed contains fish meal as the major ingredient that causes the deterioration of pond water quality leading to poor and non-uniform growth, low condition index values, low survival rate and high FCR. The present study reveals the upgradation of pond health and related success indicators of *M. rosenbergii* culture due to the use of

mangrove based plant ingredients in preparing the feed. Among the two selected floral species, *P. coarctata* generated best results. We conclude from this study that partial replacement of fish meal by mangrove associate floral species as feed ingredients can give better results in terms of water quality, survival, growth and condition state of the cultured species and it can be recommended to the local farmers for pond culture of *M. rosenbergii*.

SUMMARY OF RESEARCH

1. This work particularly emphasizes on partial replacement of fish meal (highly priced ingredient) with available mangrove floral resources for preparation of prawn feed in a sustainable manner.
2. It has availed researchers to work more on the usefulness of mangrove floral resources on the basis of their biochemical composition with reference to prawn nutrition.
3. It has also helped the researchers to know about the efficacy of mangrove based feed towards maintaining the water quality, growth performance and length-weight relationship of the culture species (*Macrobrachium rosenbergii*).

FUTURE ISSUES

From the findings, *Porteresia coarctata* based feed had the best impact on water quality as well as on the cultured species, suggesting that the particular formulation can be employed to prawn farmers of Indian Sundarbans for commercial production of freshwater prawns. This can pave the avenue of small scale cottage industry in the coastal zone of the country where mangroves are widely distributed.

DISCLOSURE STATEMENT

There is financial support for this research work from the funding agency.

ACKNOWLEDGEMENT

The authors are grateful to Department of Science and Technology, Government of India for providing financial assistance to the project [Sanction No. DST/IS-STAC/CO2-SR-59/09 (pt. II) dated 24.05.2011] and Department of Marine Science, University of Calcutta for providing its infrastructural facilities. One of the authors (Sufia Zaman) is grateful to Department of Science and Technology, Government of India for providing financial assistance (under Women Scientist Scheme- B Project, Sanction No. SSD/SS/028/2010/G dated 28.09.2011).

REFERENCES

1. Abohweyere PO, Williams AB. Length-weight relationship and condition factor of *Macrobrachium macrobrachion* in the Lagos-Lekki lagoon system, Nigeria. *Research Journal of Biological Sciences.*, 2008, 3(11), 1333-1336
2. Abrahamsson S. Dynamics of an isolated population of the crayfish *Astacus astacus* (Linne) and *Pasifastacus leniusculus* (Dana). *Svensk Naturvetenskap.*, 1966, 306-316
3. Ahmed N. Prawn and prawn fishery of East Pakistan. Government Press, Dhaka, East Pakistan, 1957, 1-31
4. AOAC. Official methods of Analysis, 13th Ed. Association of Official Analytical Chemists. Washington DC., 1980, 376-384
5. APHA. Standard methods for the examination of water and waste water. 19th Edition, American Public Health Association, 1995, USA
6. Ashmore SB, Standby RW, Moore LB, Malecha SR. Effect on growth and apparent digestibility of diets varying in gram source and protein level in *Macrobrachium rosenbergii*. *Journal of the World Mariculture Society.*, 1985, 16, 205-216
7. Banerjee K, Mitra A, Mondal K. Cost-effective and eco-friendly shrimp feed from red seaweed *Catenella repens* (Gigartinales: Rhodophyta). *Current Biotica.*, 2010, 4(1), 23-43
8. Biswas G, Sundaray JK, Thirunavukkarasu AR, Kailasam M. Length-weight relationship and variation in condition of *Chanos chanos* (Forsskal, 1775) from tide-fed brackishwater ponds of the Sunderbans- India. *Indian Journal of Geo-Marine Science.*, 2011, 40(3), 386-390
9. Boyd CE. Water Quality in Ponds for Aquaculture. Alabama Agricultural Experiment Station, Auburn University, Alabama, 1990, 482 pp
10. Cohen D, Ra'anan Z. The production of the freshwater prawn *Macrobrachium rosenbergii* in Israel: 3. Density effect of all-male tilapia hybrids on prawn yield characteristics in polyculture. *Aquaculture.*, 1983, 35, 57-71
11. D'Abramo LR, Brunson MW. Production of Freshwater Prawns in Ponds. SRAC (Southern Regional Aquaculture Center), 1996, Publication No: 484
12. D'Abramo LR, Sheen SS. Nutritional requirements, feed formulation and feeding practices for intensive culture of the freshwater prawn, *Macrobrachium rosenbergii*. *Rev Fish Sci.*, 1991, 2, 1-21
13. Daniel S. Introducing *Macrobrachium rosenbergii*. Freshwater and Marine Aquarium., 1981, 4(7), 32-34
14. De Silva SS, Anderson TA. Fish Nutrition in Aquaculture. Chapman and Hall, 1995, 2-6 Boundary Row, London SE1 8HN
15. Enin UI. Length-weight parameters and condition factor of two West African prawns. *Revue d' Hydrobiologie Tropicale.*, 1994, 27(2), 121-127
16. Fischer W, Bianchi G, Scott WB. FAO species identification sheets for fisheries purposes. Eastern Central Atlantic, fishing areas, 34, 47 (in part). Food and Agriculture Organization of the United Nations, 1981
17. Gomez KA, Gomez AA. Statistical procedures for agricultural research, 2nd Edition, 1984, John Wiley and Sons, New York, 680 pp
18. Hall MR, Van Hamm EH. The effect of different types of stress on blood glucose in giant tiger prawn *Penaeus monodon*. *Journal of the World Aquaculture Society.*, 1998, 29, 290-299
19. Hasanuzzaman AFM, Siddiqui MN, Chisty MAH. Optimum Replacement of Fishmeal with Soybean Meal in Diet for *Macrobrachium rosenbergii* (De Man 1879) Cultured in Low Saline Water. *Turkish Journal of Fisheries and Aquatic Sciences.*, 2009, 9, 17-22
20. Hossain MA, Jauncey K. Substitution of fishmeal by oilseed meals in various combinations in the diet of common carp (*Cyprinus carpio*). *Malaysian Applied Biology.*, 1990, 9, 1-12
21. Hossain MA, Paul L. Low-cost diet for monoculture of giant freshwater (*Macrobrachium rosenbergii* de Man) in Bangladesh. *Aquaculture Research.*, 2007, 38, 232-238
22. Hossain MA, Siddique MAL, Miaje MAH. Development of low-cost feed for culture of giant freshwater prawn (*Macrobrachium rosenbergii* de Man) in ponds. *Bangladesh Journal of Fisheries Research.*, 2000, 4, 127-134
23. Jayachandran KV, Joseph NI. Length-weight relationship of two palaemonid prawns *Macrobrachium idella* and *M. scabriculum*. A comparative study. *Fishery Technology.*, 1988, 25, 189-195
24. Kuris AM, Ra'anan Z, Sagi A, Cohen D. Morphometric differentiation of male Malaysian giant prawns *Macrobrachium rosenbergii*. *Journal of Crustacean Biology.*, 1987, 7, 219-237

Mondal et al.

Partial replacement of fish meal with mangrove based plant ingredients and its effect on water quality, growth performance and length-weight relationship of freshwater prawn *Macrobrachium rosenbergii*, Species, 2013, 3(8), 15-21,

© The Author(s) 2013. Open Access. This article is licensed under a [Creative Commons Attribution License 4.0 \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/).

25. Le Cren ED. The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *J Anim Ecol.*, 1951, 20, 201-219
26. Lokare KV, Anis S. Present status of prawn farming in Maharashtra. National Workshop in Aquaculture of freshwater prawns organized by C.I.F.E., Mumbai and COF, Nellore (AP), 8-9 February 2000, (Abstract No. 9)
27. Mitra A, Banerjee K, Banerjee A, Das S, Golui P, Basu S. Culture technology of freshwater prawn, *Macrobrachium rosenbergii* In: Training manual on ecofriendly and sustainable fishery. WWF-India Sundarbans landscape project, 2006, 9-34 pp
28. Mitra A, Banerjee K. Mangrove and their associates: our next door neighbour. WWF-New Delhi, 2006, 36 pp
29. Mitra A, Choudhury A. Causes of water pollution in prawn culture farms. *Journal of Indian Ocean Studies.*, 1995, 2(3), 230-235
30. Mitra A. The north-west coast of the Bay of Bengal and deltaic Sundarbans. In: Charles Sheppard (ed.) Seas at the Millennium: An Environmental Evaluation, U.K., 2000, 2, 160 pp
31. Moss RH, Brenkert AL, Malone EL. Vulnerability to Climate Change: A Quantitative Approach, Pacific Northwest National Laboratory operated by Battelle for the United States Department of Energy, Oak Ridge, TN., 2001, <http://www.pnl.gov/globalchange/pubs/vul/DOE%20VCC%20report.PDF>
32. New MB, Singholka S. Freshwater prawn farming: A manual for the culture of *Macrobrachium rosenbergii*. FAO Fisheries Technical Paper No. 225 Rev., 1985, Rome
33. New MB. Status of freshwater prawn farming: A Review. *Aquacult Res.*, 1995, 26, 1-54
34. Pérez-Osuna F. The environmental impact of shrimp aquaculture: causes, effects and mitigating alternatives. *Environmental Management.*, 2001, 28(1), 131-140
35. Pauly D. Fishbyte section Editorial. *NAGA, ICLARM Quart.*, 1993, 16, 26 pp
36. Ricker WE. Linear regression in fishery research. *J Fish Res Board Can.*, 1973, 30, 409-434
37. Sandifer PA, Smith TIJ. Freshwater prawns. In: JV Huner and EE Brown (Eds.), Crustacean and Mollusk Aquaculture in the United States. Avi Publishing, CT, USA, 1985, 63-125
38. Sarver D, Malecha S, Onizuka D. Possible sources of variability in stocking mortality in post larval *Macrobrachium rosenbergii*. In: New MB (editor). Giant Prawn Farming, Developments in Aquaculture and Fisheries Science. Elsevier Scientific Publishing, Amsterdam, The Netherlands., 1982, 10, 99-113
39. Siddiqui AQ, Al-Hafedh YS, Al-Harbi AH, Ali SA. Effects of stocking density and monosex culture of freshwater prawn *Macrobrachium rosenbergii* on growth and production in concrete tanks in Saudi-Arabia. *Journal of the World Aquaculture Society.*, 1997, 28, 106-112
40. Strickland JDH, Parsons TR. A practical handbook of seawater analysis (2nd Edition). *Journal of the Fisheries Research Board of Canada.*, 1972, 167, 311 pp
41. Tidwell JH, Coyle C, Weibel SD, Evans J. Effects and interactions of stocking density and added substrate on production and population structure of freshwater prawns *Macrobrachium rosenbergii*. *Journal of the World Aquaculture Society.*, 1999, 30, 174-179
42. Tidwell JH, Coyle SD, Sedlacek JD, Weston PA, Knight WL, Hill SJ, D'Abramo LR, Fuller MJ. Relative prawn production and benthic macroinvertebrate densities in unfed, organically fertilized, and fed pond systems. *Aquaculture.*, 1997, 149, 227-242
43. Tidwell JH, Webster CD, Sedlacek JD, Weston PA, Knight WL, Hill Jr SJ, D'Abramo LR, Daniels WH, Fuller MJ, Montafiez JL. Effects of complete and supplemental diets and organic pond fertilization on production of *Macrobrachium rosenbergii* and associated benthic macroinvertebrate populations. *Aquaculture.*, 1995, 138, 169-180
44. Walkley A, Black IA. An estimation for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science.*, 1934, 37, 29-38
45. Weidenbach RP. Dietary components of prawns reared in Hawaiian ponds. In: Proceedings of the Giant Prawn Conference, International Foundation for Science, Report 9, 1980, Bangkok
46. Weidenbach RP. Dietary components of freshwater prawns reared in Hawaiian ponds. In: M.B. New (Ed.), Giant Prawn Farming. Developments in Aquaculture and Fisheries Science., 1982, 10, 257-267
47. William HD, D'Abramo LR, Fondren MW, Durant MD. Effects of stocking density and feed on pond production characteristics and revenue of harvested freshwater prawns, *Macrobrachium rosenbergii* stocked size-graded juveniles. *Journal of the World Aquaculture Society.*, 1995, 26, 38-47
48. Wong KH, Cheung PCK. Nutritional evaluation of some subtropical red and green seaweeds. Part II. *In vitro* protein digestibility and amino acid profiles of protein concentrate. *Food Chem.*, 2001, 72, 11-17
49. Wootton RS. Fish Ecology. Printed in Great Britain by Thomson Litho Ltd., 1992, Scotland

RELATED RESOURCE

1. Balazs GH, Ross E. Effect of protein source and level on growth and performance of captive fresh water prawn, *Macrobrachium rosenbergii*. *Aquaculture.*, 1976, 7, 299-313
2. Behanan L, Mukandan M K, Korath A, Sherif PM. Formulation and evaluation of a particulated feed for *Macrobrachium rosenbergii* post-larvae. In: Freshwater prawn (Ed. by E. G. Silas). *Kerala Agriculture University*. Thrissur, India, 1992, 238-240
3. Castell JD, Kean JC, D'Abramo LR, Conklin DE. A standard reference diet for crustacean nutrition. I. Evaluation of two formulations. *Journal of World Aquaculture Society.*, 1989, 20 (3), 93 - 99
4. D'Abramo LR, Brunson MW. Production of Freshwater Prawns in Ponds. SRAC (Southern Regional Aquaculture Center), 1996, Publication No: 484
5. Davies SJ, Brown MT, Camilleri M. Preliminary assessment of the seaweed *Porphyra purpurea* in artificial diets for thick lipped grey mullet (*Chelon labrosus*). *Aquaculture.*, 1997, 152, 249-258
6. Eusebio PS, Coloso RM. Evaluation of leguminous seed meals and leaf meals as plant protein sources in diets for juvenile *Penaeus indicus*. *Israeli Journal of Aquaculture - Bamidgeh.*, 1998, 50(2), 47-54
7. Ferdose A, Hossain MB. Nutritional value of wild, cultured and frozen prawn *Macrobrachium rosenbergii* (De Man, 1879). *International Journal of Natural Sciences.*, 2011, 1(2), 52-55
8. Floreto Eric AT, Bayer RC, Brawn PB. The effects of soybean based diets, with and without amino acid supplementation, on growth and biochemical composition of juvenile American lobster, *Homarus americanus*. *Aquaculture.*, 2000, 189, 211-235
9. González-Félix ML, Silva FS, Davis DA, Samocha TM, Morris TC, Wilkenfeld J S, Perez-Velazquez M. Replacement of fish oil in plant based diets for Pacific white shrimp (*Litopenaeus vannamei*). *Aquaculture.*, 2010, 309, 152-158
10. Jayalakshmi B, Natarajan P. Growth, conversion efficiency and moulting frequency of *Macrobrachium idella* fed on different artificial feeds. *J. Aquacult. Trop.*, 1994, 9, 193-200
11. Kaushik SJ, Cravedi JP, Lalles JP, Sumpter J, Fauconneau B, Laroche M. Partial or total replacement of fish meal by soybean protein on growth, protein utilization, potential estrogenic or antigenic effects, cholesterolemia and flesh quality in rainbow trout, *Oncorhynchus mykiss*. *Aquaculture.*, 1995, 133, 257-274
12. Mitra A, Basu S, Banerjee A, Banerjee K. Culture Of Freshwater Prawn Using Feed Incorporated With Salt Marsh Grass, *Porterasia Coarctata* in Indian Sundarbans. *Seshaiyana.*, 2007, 15 (1), 16-17